Oscillations

- 1. Periodic Motion
 - 1. Sinusoidal Motion
 - 2. How do we get this kind of motion?
 - 3. Position Velocity Acceleration
 - 4. A spring with vectors
- 2. Rotational Motion Variables
 - 1. The reference circle
 - 2. The reference circle
 - 3. The math of oscillations
 - 4. Math of rotations
- 3. Frequency of Vibration
 - 1. <u>Hooke's law</u>
- 4. Energy
 - 1. <u>Energy</u>
 - 2. Elastic Potential Energy
- 5. Pendulums
 - 1. The period of the pendulum
 - 2. phase
 - 3. The physical pendulum
 - 4. <u>Time constant</u>
 - 5. Yes damping on an oscillator
 - 6. Large angles
 - 7. Large Angles
- I. Periodic Motion



Phenomena in nature can repeat. This chapter will deal with motions that recur in time.

The top graph shows the positive of an object with respect to time. The object shown starts at rest at position +A, then moves at constant velocity towards position -A, then pauses, then returns to position +A and starts the motion over. This is an example of **periodic motion**.

The lower graph shows an object undergoing a similar motion, except for some small differences. Instead of stopping for a long time, and then moving at constant velocity, this object slows down, pauses for an instant, then speeds up in the opposite direction, only to start slowing down again. This type of motion can be described by a sinusoidal function. And will be the main type of motion we look at.

1.1 Sinusoidal Motion

The simplest case is a sinusoidal motion. Here the graph has a sine or cosine shape.



This type of motion is called "simple harmonic motion"

1.2 How do we get this kind of motion?

Simple harmonic motion will result from a restoring force, like Hooke's Law:



This is an example of a **linear restoring force**: the net force is towards the equilibrium position and is proportional to the distance from equilibrium.

 $m{k}$ is a constant known as the 'spring constant'



The force is proportional to the displacement.



```
Quick Question 1
```

What is the velocity and force at point P? (+A is to the right)







Here is the acceleration of a mass with respect to time. At which point is the speed of the mass zero? (Answer E if never at rest)



At which point(s) is the velocity of the mass the most negative ?

2. Rotational Motion Variables



Linear Velocity, v, referred to motion in x or y directions (or a combination).

Angular velocity, $\boldsymbol{\omega}$, refers to the motion of an object around a center point.

$$\overline{\omega} = rac{\Delta heta}{\Delta t}$$

$$\overline{\omega} = rac{ heta_f - heta_i}{t_f - t_i}$$

This will be easier in the long run, although it takes some getting used to.

2.4 The reference circle

Imagine shining a light at a rotating table.

The shadow cast by the rotating object would create a sinusoidal curve in time.

This allows us to always correlate simple harmonic motion with the displacement of an object around a circle.

2.5 The reference circle

Notes for PHYS20300 - Oscillations J. Hedberg, 2025



$$x = A \cos \theta$$

But, since the angular velocity $\boldsymbol{\omega}$ times the time, \boldsymbol{t} equals the angular displacement:

$$\omega t = heta$$

we can also write:

$x = A \cos \omega t$

2.6 The math of oscillations

Frequency and Period are used to describe a repeating motion:

$$f=rac{1}{T}$$
 $T=rac{1}{f}$

So far, we have usually talked about period, that is how long for one oscillation. Frequency is probably more useful for many applications though.

Frequency is the number of cycles per second.

Frequency: 1 Hertz is the unit of frequency:

2.7 Math of rotations

We'll need to be able to express the various characteristics of an oscillating system:

The angular velocity, $\boldsymbol{\omega}$ was "radians per second"

Thus, one rotation is 2π radians, so we can see that the frequency is:

$$\omega = 2\pi f$$

Likewise for the period:

$$\omega = rac{2\pi}{T}$$

Notes for PHYS20300 - Oscillations J. Hedberg, 2025



The displacement of an object in SHM is

 $x = A \cos \omega t$

The velocity can be obtained from a little geometry, or calculus:

 $v = -A\omega\sin\omega t$

And lastly, the acceleration is given by:



If an object has a velocity given by: $v(t) = -B\sin\omega t$ what will its maximum speed be?

3. Frequency of Vibration

These springs mass systems have different frequencies of vibration

Starting with Hooke's Law for the force on a spring

F = -kx

Since Newton's second law says that F = ma, we can then write:

$$-k(A\cos\omega t) = m(-A\omega^2\cos\omega t)$$

Let's solve this for $\boldsymbol{\omega}$:

$$\omega = \sqrt{rac{k}{m}}$$

3.8 Hooke's law

The spring constant ${m k}$ tells us 'how stiff the spring is'

The units of the spring constant are Newtons per meter (N/m)

Below is plotted the position vs. time for three ball & spring systems. Which one has the smallest spring constant ?





Notes for PHYS20300 - Oscillations J. Hedberg, 2025



3.10 Elastic Potential Energy

The elastic potential energy of a spring mass system tells us how much energy is stored in the spring.

$$PE = rac{1}{2}kx^2$$

When the displacement is zero from the equilibrium, then no potential energy exists in the system.

We can see the transformation from potential to kinetic and back again in a simple harmonic oscillator



The total energy remains constant:

$$E=rac{1}{2}mv^2+rac{1}{2}kx^2$$

mass has zero velocity at endpoints (x = A or -A, the turning points); all energy is potential at those points

mass has greatest speed at equilibrium (x = 0); all energy is kinetic



Thus, let's solve for v_{\max}

 $v_{ ext{max}} = \sqrt{rac{k}{m}} A$

But,

$$v_{
m max} = A \omega$$

Thus,

$$\omega = \sqrt{rac{k}{m}}$$
 $f = rac{1}{2\pi} \sqrt{rac{k}{m}} ~~{
m or}~~T = 2\pi \sqrt{rac{m}{k}}$

The frequency and period are determined by the physical properties of the oscillator.

All that matters is the mass of the object and the spring constant.

The frequency and period do not depend on the amplitude A.

Small and large oscillations have the same frequency.

Whenever we have this condition, of energy trading off between kinetic and potential, we'll get an oscillatory behaviour.

Some are easier to describe than others.

4. Pendulums

A simple pendulum consists of a mass m attached to a frictionless pivot by a cable of length L.





For small angles, the tangential force on the pendulum bob will be given by:

$$F_{
m tang} = -mg\sin heta$$

When θ is very small ($\theta < .2$ radians), $\sin \theta = \theta$.

$$F_{ ext{tang}} = -rac{mg}{L}s$$

This is essentially a restoring force, just like we had for the mass/spring system.

$$F=-kx \;\;\Rightarrow\;\; F=-rac{mg}{L}s$$

Now $k
ightarrow rac{mg}{L}$, thus we could replace our frequencies of the spring mass systems with:

$$f_{
m spring-mass} = rac{1}{2\pi} \sqrt{rac{k}{m}} \; \Rightarrow \; f_{
m pendulum} = rac{1}{2\pi} \sqrt{rac{g}{L}}$$

4.11 The period of the pendulum

Since $f_f = \frac{1}{2\pi} \sqrt{\frac{g}{L}}$, we can easily write the period of a pendulum. This is how long it takes to do one oscillation:

$$T_{
m pend}=2\pi\sqrt{rac{L}{g}}$$

This is only true for *small oscillations*. But, we can still do a lot with small oscillations.

When the pendulum is at the lowest point, what can you say about its acceleration? (Consider all possible accelerations!)

Quick Question 6

A grandfather clock, which uses a pendulum to keep accurate time, is adjusted at sea level. The clock is then taken to an altitude of several kilometers. How will the clock behave in its new location?

- a) The clock will run slow.
- b) The clock will run fast.

c) The clock will run the same as it did at sea level.

Damping

An exponential decay function:

$$x(t) = A e^{-t/ au}$$

Now, au is not torque, but will be called the **time constant** for this decaying curve.

$$x(t= au)=Ae^{-1}$$

$$rac{A}{e}pprox .37A$$

The time constant describes how quickly the exponential function approaches zero.

4.12 Time constant

Different time constants will effect the graphs in this way.

$$au=.5$$
 $au=1$
 $au=2$

The time constant, au, can be defined: "The time where the value gets to 1/e of the initial value, or about a third (.37)"

No damping on an oscillator

An oscillator that is not damped, and loses no energy will have a motion like this:

$$x(t) = \sin\left(rac{2\pi t}{T}
ight)$$

4.13 Yes damping on an oscillator

However, if we damp an oscillator, then we'll see a decay in the amplitude of the oscillations.

$$x(t)=e^{rac{-t}{ au}}\sin\!\left(rac{2\pi t}{T}
ight)$$

These oscillation have the same period, and frequency, and A. However, the au which dictates the damping is different.

$$x(t) = e^{rac{-t}{ au}} \sin\!\left(rac{2\pi t}{T}
ight)$$

Here, the different time constants are given by:

$$au = \{.5, 1, 2\}$$

Damping



The damping leads to a different $\omega
ightarrow \omega'$

$$\omega'=\sqrt{rac{k}{m}-rac{b^2}{4m^2}}$$

Driving a resonator

All the oscillators in the real world will have some kind of damping force: friction, drag, thermal losses. If we want this pendulum to keep swinging, we need to add energy to account of the lost energy.

In order to keep things oscillating, we have to drive the oscillations. A driven oscillation has an applied force which also oscillates at some frequency. There will be a natural frequency of oscillation. In the case of the pendulum, that's

$$f_0=rac{1}{2\pi}\sqrt{rac{g}{L}}$$

If the oscillator is driven with a matching frequency, then we have the maximum amplitude of displacement.

(Nearly everything has a natural resonance frequency - buildings, bridges, wine glasses)

4.14 Large angles



Our previous analysis relied on the approximation that $\sin(\theta) \approx \theta$. What if we have larger values of θ where this is not true?

4.15 Large Angles

If instead we have to solve this equation:

$$lpha = rac{d^2 heta}{dt^2} = -rac{mg}{L}{\sin heta}$$

We'll need fancier math.

We'll end up with something that looks like this:

$$T = 2\pi \sqrt{\frac{L}{g}} \left(1 + \frac{1}{16}\theta_0^2 + \frac{11}{3072}\theta_0^4 + \frac{173}{737280}\theta_0^6 + \frac{22931}{1321205760}\theta_0^8 + \frac{1319183}{951268147200}\theta_0^{10} + \dots \right)$$
(1)

Example Problem #1:

A 2 kg mass is attached to a spring and stretched 4 cm in the positive x direction. At t = 0, the mass is released and the system begins to oscillate at a frequency of 2 Hertz.

- 1. Write equations that describe the position, velocity, and acceleration of the mass.
- 2. Sketch 3 plots that show these three quantities as functions of time.
- 3. Compile a table of important values: f, T, ω , A, v_{max} , total energy, k.

Example Problem #2:



1. What is the equilibrium length of the spring?

- 2. Where are the minimum and maximum extensions? (i.e. the turning points)
- 3. What is the particle's maximum kinetic energy?
- 4. If the total energy of the particle is doubled, where will the turning points be?

Example Problem #3:

A ball of mass m oscillates on a spring with spring constant k = 200N/m. The ball's position is given by: $x = .350m \times \cos(15.0t)$ with t measured in seconds.

- 1. Find the amplitude of the motion
- 2. Find the frequency of the motion
- 3. What is the mass
- 4. What is the total energy
- 5. That is the maximum speed of the ball?

Example Problem #4:

On a cruise ship in the ocean, the waves make the boat rock up and down. A passenger experiences a vertical motion of amplitude 1 m with a period of 15 s. (a) What is the maximum acceleration this passenger feels? (b) What fraction of g is this?

Example Problem #5:

A 200 g air-track glider is attached to a spring. The glider is pushed 10 cm against the spring, then released. 10 oscillations are found to take 12 seconds. What is the spring constant of this spring?