

Stellar Interiors

1. [Hydrostatic Equilibrium](#)
 1. [Pressure in the sun](#)
 2. [Pressure-radius](#)
2. [Mass continuity](#)
3. [Stellar Energy Sources](#)
 1. [Where does it come from?](#)
4. [Intro to Nuclear Reactions](#)
 1. [Fission](#)
 2. [Fusion](#)
 3. [Fundamental Forces](#)
 4. [Binding Energies](#)
 5. [Bringing them together](#)
 6. [Energy / Tunneling](#)
 7. [Cross Section, basics](#)
 8. [Cross Section](#)
 9. [p-p chain](#)
 10. [CNO Cycle](#)
5. [Luminosity Gradient](#)
6. [Energy Transport](#)
 1. [Radiative](#)
 2. [Convective Transport](#)
7. [Stellar Model Building](#)





Figure shows the basic structure of a star. (It is not scale!) The light we see comes from photons in the stellar atmosphere.

The basic structure of a star.

I. Hydrostatic Equilibrium



Let's imagine a small cylindrical volume of gas that is located in a larger region. There will be three forces acting on it: F_{grav} is the force of gravity, and two forces from the pressure, one from the pressure above: $P + \Delta P$ and one from pressure below: P . The net force from the pressure will be :

$$F_{\text{pres}} = A[P - (P + \Delta P)] \quad (1)$$

A small volume of gas has three force vectors acting in the vertical directions: gravity, pressure from below and pressure from above.

where A is the area (whose surface normal points in the y direction) of the gas volume. The gravitation force will be given by:

$$F_{\text{grav}} = -\frac{GM_r \rho A \Delta r}{r^2} \quad (2)$$

where M_r is the mass enclosed by a sphere of radius r centered about the earth's center.

If this volume of gas is in equilibrium, then the pressure forces and the gravitational forces will be equal:

$$A[P - (P + \Delta P)] - \frac{GM_r \rho A \Delta r}{r^2} = 0 \quad (3)$$

Solving for ΔP :

$$\Delta P = -\frac{GM_r \rho}{r^2} \Delta r \quad (4)$$

If we consider this cylinder to be infinitely thin: $\Delta r \rightarrow 0$, then we have a differential form:

$$\frac{dP}{dr} = -\frac{GM_r \rho}{r^2} \quad (5)$$

Eq. (5) is called the **equation of hydrostatic equilibrium** (for a spherical symmetric body).

To apply this to a star, we should acknowledge the limiting assumptions we made:

1. The star is spherical
2. The star is not rotating
3. The star is not expanding or contracting
4. Gravity and Pressure gradients are the only forces acting.

$$\frac{dP}{dr} = -\frac{GM_r \rho}{r^2} \quad (6)$$

Example Problem
#1:

1.1 Pressure in the sun

Find a rough estimate for the pressure at the center of the sun.

$$\frac{dP}{dr} = -\frac{GM_r \rho}{r^2} \quad (7)$$

can be expressed as a difference equation:

$$\frac{dP}{dr} \sim \frac{P_s - P_c}{R_s - 0} \sim -\frac{P_c}{R_\odot} \quad (8)$$

1.2 Pressure-radius

A better way to write this would be to include the dependencies:

$$\frac{dP}{dr} = -\frac{GM(r)\rho(r)}{r^2} \quad (9)$$

We can see that there are 3 quantities that depend on r : Pressure (P), Mass (M) and Density (ρ). This makes a unique solution not possible. We need more equations!

2. Mass continuity

$$\frac{dM}{dr} = 4\pi r^2 \rho(r) \quad (10)$$

The **equation of mass continuity** relates density and radial position and total mass. In short it says that the total mass is just the sum of all the infinitesimal spherical shells (i.e. onion skins)

3. Stellar Energy Sources

3.3 Where does it come from?

The first suggestion to explain where stars got the energy was to look at the gravitational potential. Let's say we have two particles, then the gravitational potential energy between them is:

$$U = -G \frac{Mm}{r} \quad (11)$$

If the distance r between these particles decreases, then the potential energy must get more negative, and we would expect to find energy in other forms, kinetic for example.

How much energy is there in gravity?

Of course there are more than two particles in a star, so we'll have to expand that equation a bit. If a point mass dm is located outside a spherically symmetric mass M_r , then the force on that point mass will be directed to the center and will have a magnitude:

$$dF_{g,i} = G \frac{M_r dm_i}{r^2}$$

The potential energy of the point mass is thus:

$$dU_{g,i} = -G \frac{M_r dm_i}{r} \quad (12)$$

Now we'll consider a shell of thickness dr and mass dm :

$$dm = 4\pi r^2 \rho dr$$

where ρ is the density and $4\pi r^2 dr$ is the volume of the shell. Thus, eq (12) becomes

$$dU_g = -G \frac{M_r 4\pi r^2 \rho}{r} dr$$

We can integrate over the radius of the star R like this:

$$U_g = -4\pi G \int_0^R M_r \rho r dr \quad (13)$$

Of course we probably don't know ρ very well at this point, but we can guess based on it's average value:

$$\rho \sim \bar{\rho} = \frac{M}{\frac{4}{3}\pi R^3}$$

with M being the total mass of the star. Also:

$$M_r \sim \frac{4}{3} \pi r^3 \bar{\rho}$$

Putting this into eq. (13), we get:

$$U_g \sim \frac{16\pi^2}{15} G \bar{\rho}^2 R^5 \sim -\frac{3}{5} \frac{GM^2}{R} \quad (14)$$

Applying the virial theorem, we have:

$$E \sim -\frac{3}{10} \frac{GM^2}{R} \quad (15)$$


If we plug some sun values into this:

$$\frac{3}{10} \frac{GM_{\odot}^2}{R_{\odot}} \approx 1.1 \times 10^{41} \text{ J}$$

Now if the sun luminosity was constant at today's value: $3.84 \times 10^{26} \text{ W}$, then this would mean there is enough energy to last for about 1×10^7 years (about 10 million years). This is notably too short given other observations from geology that suggest the earth and moon and things have been around for billions of years. There must be another source of energy contributing to the star's luminosity.

4. Intro to Nuclear Reactions


4.4 Fission



When an atom breaks apart. Power plants...

<http://www.nuclear-power.net/nuclear-power/fission/>

4.5 Fusion



When you smash two atoms together. Suns...

<http://www.nuclear-power.net/nuclear-power/nuclear-fusion/>

Take four Hydrogen atoms (1 proton & 1 electron), each have a mass of 1.00782503214 u. Together, they therefore have a combined mass of 4.03130013 u. One Helium-4 atom has a mass of 4.002603 u. This is obtained by adding the mass of the neutrons and the protons together. Thus, let's say we were able to mash 4 Hydrogen atoms together and make a Helium. Now there would be a change in mass between the initial and final constitutions:


$$\Delta m = 4.03130013 - 4.002603 = 0.028697 \text{ u}$$

Since $E = mc^2$, we can see that this amount of mass is associated with an energy of 26.731 MeV. This is essentially the amount of energy released in the formation of a Helium nucleus, aka the **binding energy**. (This is also the amount of energy that would be required to take apart a Helium nucleus.) The trick is to figure out exactly how we can combine 4 Hydrogen atoms to make a Helium. (that's the study of High energy physics earlier last century)

4.6 Fundamental Forces


Force	Relative Strength	Distance	What does?
Weak	10^{33}	10^{-18}	Radioactive Decay
Strong	10^{38}	10^{-15}	Keeps atomic nuclei together
Gravity	1	∞	Brings everything together
Electromagnetic	10^{36}	∞	Chemistry, charges, etc

Scales →



The Isotopes of Hydrogen

4.7 Binding Energies




Carroll & Ostlie, figure 10-9.

4.8 Bringing them together

One impediment is the Coulomb repulsion between two protons, both of which are positively charged.

As the separation distance between two charged (alike) particles decreases, the force of repulsion increases. The Coulomb potential between two protons located 1 fm apart would be:

$$\begin{aligned} &\approx \frac{(1.60 \times 10^{-19} \text{ C})^2}{4\pi(8.8 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1})(1.60 \times 10^{-19} \text{ J eV}^{-1})(10^{-15} \text{ m})} \\ &\approx 1.4 \times 10^6 \text{ eV} \\ &\approx 1.4 \text{ MeV} \end{aligned}$$



If two protons wanted to get closer than 1 fm, they would need to have a kinetic energy greater than this value to overcome the potential barrier. The energy of a proton at the center of the sun is about:

$$\langle E \rangle = \frac{3kT_c}{2} \approx \frac{3(1.38 \times 10^{-23} \text{ J K}^{-1})(1.47 \times 10^7 \text{ K})}{2(1.6 \times 10^{-19} \text{ J eV}^{-1})} \approx 2 \text{ keV} \quad (16)$$

This would be insufficient energy to get over the barrier. This is where quantum tunneling comes in.

The plots of the Coulomb force (top) and the Coulomb potential (bottom).

Tunneling

Criteria for tunneling: a) the distance between protons should be comparable to the de Broglie wavelength:

$$\lambda_{DB} = \frac{h}{p} = 7 \times 10^{-13} \text{ m} \quad (17)$$

and b) the kinetic energy (E) of the proton needs to be comparable to the electrostatic potential energy at that separation.


$$U = \frac{e^2}{4\pi\epsilon_0} \frac{1}{\lambda_{DB}} = \frac{e^2}{4\pi\epsilon_0} \frac{(2m_p E)^{1/2}}{h} \approx E \quad (18)$$

Solving for the kinetic energy:

$$E \approx \left(\frac{e^2}{4\pi\epsilon_0} \right)^2 \frac{2m_p}{h^2} \approx \frac{1}{2\pi^2} \alpha^2 m_p c^2 \approx 3 \text{ keV} \quad (19)$$

Since this energy is comparable to the kinetic energy of a proton in the core of the sun, we can expect tunneling through the Coulomb barrier to be possible. The next step is to figure out just *how likely* such events are.

4.9 Energy / Tunneling




A qualitative graph showing the what happens when two protons get close enough for the strong force to dominate.

4.10 Cross Section, basics

We can say each apple has a cross-section of:

$$N_{\text{apples}} = 13$$



Apples in a tree



If we shoot 200 arrows into the tree, how many apples will we hit?

$$\sigma = \pi r^2$$

$$N_{\text{hit}} = N_{\text{inc}} * n_{\text{tar}} * \sigma$$

where $n_{\text{tar}} = \frac{\text{number of apples}}{\text{area of tree}}$ and $n_{\text{inc}} = \# \text{ incident arrows} = 200$ If $r = 0.1$ and $R = 5.0$ then,

$$N_{\text{hit}} = 200 \times \frac{13}{\pi 5.0^2} \times \pi 0.1^2$$

Or, we can divide the number of apples by a Δt to get a *rate*, then we would have a rate of apple hits:


$$\frac{N_{\text{hit}}}{\Delta t} = \frac{N_{\text{inc}}}{\Delta t} * n_{\text{tar}} * \sigma$$

4.1.1 Cross Section

The **cross section** gives a probability of interaction. It is the number of reactions per target nucleus per time divided by the flux of incident particles.

$$\sigma(E) \equiv \frac{\text{number of reactions/nucleus/time}}{\text{number of incident particles/area/time}}$$

Cross Section



Imagine a cylinder of length $ds = v(E)dt$ and area $\sigma(E)$.

n : just a particle

n_i : an incident particle

$n_i dE$: # of incident particles having energies between E and $E + dE$ per unit volume

dN_E : # of particles that "hit the wall" / time or # of reactions / time

$n_E dE$: # of particles with an energy between E and $E + dE$, (might not be incident)

If a particle hits the wall, then there is a reaction. This is an abstraction of the nuclear collision process. (there is no cylinder, no wall...)

We have a target particle (x) and an incident particle (i). If the incident particle strikes the area $\sigma(E)$, then there will be a nuclear reaction.

Take the number of incident particles per unit volume having energy between E and $E + dE$ to be: $n_i dE$. (This is like a density) To find the number of such particles inside the cylinder:

$$dN_E = \underbrace{n_i dE}_{\text{density}} \underbrace{\sigma(E) v(E) dt}_{\text{volume}}$$

The number of particles with the appropriate kinetic energy is some fraction of the whole:

$$n_i dE = \frac{\int_0^\infty n_i dE}{\int_0^\infty n_E dE} n_E dE = \frac{n_i}{n} n_E dE$$

where $n_E dE$ is given by the Maxwell-Boltzmann distribution

$$n_E dE = \frac{2n}{\sqrt{\pi}} \left(\frac{1}{kT} \right)^{3/2} E^{1/2} e^{-\frac{E}{kT}} dE \quad (20)$$

The number of reactions per target nucleus per time dt having energies between E and $E + dE$ is

$$\frac{\text{reactions per nucleus}}{\text{time interval}} = \frac{dN_E}{dt} = \sigma(E)v(E) \frac{n_i}{n} n_E dE$$

Lastly, if there are n_x targets per unit volume, then the total number of reactions per unit volume per unit time, given all possible energies is:

$$r_{ix} = \int_0^\infty n_x n_i \sigma(E) v(E) \frac{n_E}{n} dE \quad (21)$$

$\sigma(E)$

But, what exactly is $\sigma(E)$?

Should be in the order of the de Broglie λ

$$\sigma(E) \propto \pi \lambda_{DB}^2 \propto \pi \left(\frac{h}{p} \right)^2 \propto \frac{1}{E} \quad (22)$$

(Since $K = E = \mu_m v^2 / 2 = p^2 / 2\mu_m$)

Also, for a barrier U and particle with energy E , the probability for tunneling through will go as:

$$\sigma(E) \propto e^{-2\pi^2 U/E} \quad (23)$$

In this case the U is the Coulomb potential: $U_c = \frac{Z_1 Z_2 e^2}{4\pi\epsilon_0 r}$ and E is just the kinetic energy: $E = \mu_m v^2 / 2$. We can assume the distance r is basically the de Broglie wavelength: $r \sim \lambda = h/p$.

In which case we obtain:

$$\sigma(E) \propto e^{-bE^{-1/2}} \quad (24)$$

where b is given by


$$b \equiv \frac{\pi \mu_m^{1/2} Z_1 Z_2 e^2}{2^{1/2} \epsilon_0 h}$$

We can combine eq (22) and (24) and add another function $S(E)$ (that doesn't change much) to get a functional form for $\sigma(E)$:

$$\sigma(E) = \frac{S(E)}{E} e^{-bE^{-1/2}} \quad (25)$$

Finally, we can write a reaction rate as:

$$r_{ix} = \left(\frac{2}{kT} \right)^{3/2} \frac{n_i n_x}{(\mu_m \pi)^{1/2}} \int_0^\infty S(E) e^{-bE^{-1/2}} e^{-E/kT} dE \quad (26)$$



The peak will be at


$$E_0 = \left(\frac{bkT}{2} \right)^{2/3} \quad (27)$$

The likelihood that a nuclear reaction will occur.




The 7th Solvay Conference

<https://catalog.archives.gov/id/7665680>



4.12 p-p chain



The full p-p chain

e^+	positron
ν_e	electron neutrino
γ	gamma ray

One way to convert Hydrogen to Helium is through the proton-proton chain (PP1). The overall reaction combines 4 protons to create 1 Helium-4 plus some other stuff.

The first step in this reaction take two protons and produces a deuterium, a positron, and an electron neutrino.

$${}^1_1\text{H} + {}^1_1\text{H} \rightarrow {}^2_1\text{H} + e^+ + \nu_e$$

(28)


Then the deuterium combines with another proton to make a Helium-3 atom plus a gamma ray.



Lastly, two Helium-3s combine to create Helium-4 and 2 protons




4.13 CNO Cycle



In stars that more than $1.3M_{\odot}$, a different nuclear fusion process will dominate.

This is the **CNO cycle**. It's more complicated, but the basic idea is that Carbon, Nitrogen, and Oxygen act as catalysts to speed the fusion of hydrogen. The net result is:



The CNO cycle

5. Luminosity Gradient

$$dL = \epsilon dm \quad (32)$$

Here, ϵ is the total energy released per kilogram per second by all the star stuff: gravity and nuclear reactions together.

For a spherically symmetric star, the mass of a thin shell of thickness dr will be:

$$dm = dM_r = \rho dV = 4\pi r^2 \rho dr \quad (33)$$

which leads to:

$$\frac{dL_r}{dr} = 4\pi r^2 \rho \epsilon \quad (34)$$

6. Energy Transport

6.14 Radiative

Radiation Pressure

Energy density of Blackbody radiation:

$$u_\nu d\nu = \frac{4\pi}{c} B_\nu d\nu = \frac{8\pi h \nu^3 / c^3}{e^{h\nu/kT} - 1} d\nu \quad (35)$$

We can integrate this over all wavelengths/frequencies

$$u = \int_0^\infty u_\lambda d\lambda = \int_0^\infty u_\nu d\nu$$


Thus we obtain:

$$u = \frac{4\pi}{c} \int_0^\infty B_\lambda(T) d\lambda = \frac{4\sigma T^4}{c} = aT^4 \quad (36)$$

(a is the radiation constant given by: $7.565767 \times 10^{-16} \text{ J m}^{-3} \text{ K}^{-4}$)

The **radiation pressure** can be shown to be one-third of the energy density:

$$P_{\text{rad}} = \frac{aT^4}{3} \quad (37)$$



A thin spherical shell centered on the star center. Let $dr \ll r$

The radiation pressure on the inner surface:

$$P_{\text{rad}}(r) = \frac{aT^4}{3} \quad (38)$$

The radiation pressure on the outer surface:

$$P_{\text{rad}}(r + dr) = \frac{a(T + dT)^4}{3} = \frac{a}{3}T^4 \left(1 + \frac{dT}{T}\right)^4 \approx \frac{a}{3}T^4 \left(1 + 4\frac{dT}{T}\right) \quad (39)$$

(assuming $dT/T \ll 1$)

Thus, the net radiation force will be given by:

$$F_{\text{rad}} = [P_{\text{rad}}(r) - P_{\text{rad}}(r + dr)]4\pi r^2 \quad (40)$$

which using eqs. (38) and (39), we can get:

$$F_{\text{rad}} \approx -\frac{a}{3}4T^4 \frac{dT}{T} 4\pi r^2 = -\frac{16\pi}{3}ar^2T^3 dT \quad (41)$$

The optical depth is given by:

$$d\tau = -\rho(r)\kappa(r)dr \quad (42)$$

where κ is the opacity. The probability that a photon will be absorbed while passing through the shell is $dP \approx d\tau$.

The rate at which photons carry energy through the shell is the luminosity, $L(r)$. Momentum is carried through the shell is just $L(r)/c$, since $p = E/c$.

The force on the shell, i.e. the rate at which photon momentum is transferred to the shell will be

$$F_{\text{rad}} = \frac{L(r)}{c}d\tau = -\frac{L(r)}{c}\rho(r)\kappa(r)dr \quad (43)$$

Combining eq. (43) equation with eq. (41), we get:

$$-\frac{16\pi}{3}ar^2T^3 dT = -\frac{L(r)}{c}\rho(r)\kappa(r)dr \quad (44)$$

This can be rearranged to yield the **equation of radiative energy transport**:

$$\frac{dT}{dr} = -\frac{3\rho(r)\kappa(r)L(r)}{16\pi ac T(r)^3 r^2} \quad (45)$$


This equation related the luminosity to the temperature gradient of a star.

6.15 Convective Transport

This is the **equation of convective energy transport**.

$$\frac{dT}{dr} = \left(1 - \frac{1}{\gamma}\right) \frac{T(r)}{P(r)} \frac{dP}{dr} \quad (46)$$

It shows that the temperature gradient is proportional to the pressure gradient. (γ is the **adiabatic index** and depends on the gas in question.)



The different transport modes are separated into regions.

7. Stellar Model Building

$$\frac{dP}{dr} = -\frac{GM_r \rho}{r^2}$$

$$\frac{dM}{dr} = 4\pi r^2 \rho(r)$$

$$\frac{dL}{dr} = 4\pi r^2 \rho(r) \epsilon(r)$$

$$\frac{dT}{dr} = -\frac{3\rho(r)\kappa(r)L(r)}{16\pi ac T(r)^3 r^2} \quad [\text{for radiative}]$$

$$= \left(1 - \frac{1}{\gamma}\right) \frac{T(r)}{P(r)} \frac{dP}{dr} \quad [\text{for convective}]$$

These are the basic equations that show the relationship between the main attributes of a star. They are 4 differential equations that relate: pressure, mass, density, temperature, energy generation, luminosity, and the radius of the star.

Boundary Conditions

In order to solve these equations, we need boundary conditions: At the center we should expect:

$$\left. \begin{array}{l} M_r \rightarrow 0 \\ L_r \rightarrow 0 \end{array} \right\} \text{ as } r \rightarrow 0$$

and at the surface of the star:

$$\left. \begin{array}{l} T \rightarrow 0 \\ P \rightarrow 0 \\ \rho \rightarrow 0 \end{array} \right\} \text{as } r \rightarrow R_*$$