

Cosmology


1. [Olbers Paradox](#)
2. [Distances](#)
 1. [Standard Candles](#)
 2. [Hubble's Velocity vs. Distance curve](#)
 3. [Hubble Parameter](#)
 4. [Light](#)
3. [Expansion](#)
 1. [Hubble Flow](#)
4. [Earlier times](#)
 1. [Big Bang Implications](#)
5. [Cosmic Microwave Background](#)
 1. [Isotropy/Anisotropy](#)
 2. [Olber's Paradox Resolution](#)



A forest in Oregon

An early question about the nature of the universe is known as the Olbers's Paradox: why is it dark at night?

1. Olbers Paradox



Similarly, if we look in every direction, there should be a star there, ... eventually.

The assumptions:

1. The Universe is homogenous
2. The Universe is infinitely large
3. The Universe is infinitely old

If the universe is infinite, then there should be a star everywhere we look - why is it dark out?

2. Distances

Many different methods for measuring distances

Radar

How long for light to travel.

Example: Lunar Laser Ranging RetroReflector (LRRR) on the moon. Measures distance to solar system objects.



Goddard's Laser Ranging Facility directing a laser (green beam)


Credit: Tom Zagwodzki/Goddard Space Flight Center



Apache Point Observatory 3.5m telescope

Credit: Apache Point Observatory 3.5m telescope running APOLLO lunar laser ranging experiment taken by Dan Long

Parallax



Bessel's heliometer

By <http://www.klima-luft.de/steinicke/ngcic/persons/auwers.htm>
- <http://www.klima-luft.de/steinicke/ngcic/persons/auwers.htm>,
Public Domain,
<https://commons.wikimedia.org/w/index.php?curid=8535990>

On earth: about 100 parsecs max distance.

HST Max distance 3,000 parsecs (10,000 ly)

Gaia (10,000 parsecs with 20% error)

Milky way is at least 30,000 parsecs in diameter.

(Andromeda is about 780,000 parsecs away)

2.1 Standard Candles

The flux (measured brightness) of an object depends on two things: how far away it is and its luminosity:

$$F = \frac{L}{4\pi r^2} \tag{1}$$

where L is the luminosity and r is the distance between source and observer.

If we have a known value for L , then we can calculate r based on the measured flux.

Some well understood physical processes can help lock in a value for L

Cepheids - Variable stars


Period of brightness fluctuations is related to apparent magnitude, m .

If we know the distance to some nearby ones (from parallax), we can calibrate the relationship.

Max: 30,000,000 parsecs (~100 million ly)

Useful for objects in this galaxy, and a few nearby ones.


Type 1A Supernova




X-ray: [NASA/CXC/Rutgers/G.Cassam-Chena, J.Hughes et al.](#); Radio: [NRAO/AUI/NSF/GBT/VLA/Dyer, Maddalena & Cornwell](#); Optical: [Middlebury College/F.Winkler](#), [NOAO/AURA/NSF/CTIO Schmidt & DSS](#)

A white dwarf in a binary system can gain mass from the companion until it reaches a limit, producing an explosion. This is a supernova. The luminosity of the resulting explosion can be predicted.


Max: 1 billion ly



[STScI](#)




SN 1054 - The Guest Star



Crab Nebula, Via Hubble Space Telescope

NASA, ESA, J. Hester and A. Loll (Arizona State University)



Light Curve from Supernovas

<https://newscenter.lbl.gov/2014/03/03/standard-candle-supernovae/>

This scientific visualization shows the development of Supernova 1987A, from the initial blast observed three decades ago to the luminous ring of material we see today.

<https://hubblesite.org/contents/media/videos/939-Video>

SC22: Visualizing the Supernova Explosion of a 25

SC Conference Series




Watch on



The Vera Rubin Telescope: This survey will observe the entire visible southern sky every few nights over the course of a decade, capturing about 1000 images of the sky every night and giving us a new view of our evolving Universe

<https://noirlab.edu/public/images/iotw2207a/>


Redshift



Earlier 20th century observations of galactic spectra indicated red-shifted spectral lines for most. Vesto Slipher did the first measurements of redshift in galactic spectra. This suggested they were moving away from us.


Slipher's 1917 data on galaxies apparent velocities. (nearly all are moving away from us)

[Slipher, galaxies, and cosmological velocity fields](#), John A. Peacock



Hubble's spectra from other galaxies.


Hubble was the first to correlate the redshifts to distance using Cepheid Variable stars outside the Milky Way.



Hubble - The Realm of the Nebulae

2.2 Hubble's Velocity vs. Distance curve

<https://www.quantamagazine.org/the-webb-telescope-further-deepens-the-biggest-controversy-in-cosmology-20240813/>



A Hubble diagram using Type 1A supernova. (Hubble's original plot is contained in the small red square in the bottom left.)

[10.1073/pnas.2536799100](https://doi.org/10.1073/pnas.2536799100)

2.4 Light

Define redshift as change in wavelength w.r.t wavelength at time of emission, e .

$$z \equiv \frac{\lambda_0 - \lambda_e}{\lambda_e} = \frac{\Delta\lambda}{\lambda_e} \quad (3)$$

3. Expansion

If everything seems to be moving away from everything else, then one possibility is that space is just getting bigger.

$$\mathbf{r}(t) = a(t)\mathbf{r}_0 \quad (4)$$

where \mathbf{r}_0 is the separation at the current time, and $a(t)$ is a dimensionless function known as the scale factor.

Wavelength - scale factor relation

$$\frac{\lambda_e}{a(t_e)} = \frac{\lambda_0}{a(t_0)} \quad (5)$$

Thus,

$$1 + z = \frac{\lambda_e}{\lambda_0} = \frac{a(t_0)}{a(t_e)} = \frac{1}{a(t_e)} \quad (6)$$

Thus, observing a quasar with $z = 6.4$ means we are looking at it when the universe was

$$a(t_e) = \frac{1}{7.4} = 0.135 \quad (7)$$

times smaller.

3.5 Hubble Flow

The motion of galaxies as part of the expansion is called the **Hubble Flow**

This can be distinguished from the **peculiar velocity**, which would be for example, the motion of the Milky Way towards Andromeda.

4. Earlier times

If we run the 'film' backwards, then earlier in time, everything would have been much closer together.

Two options: either things would be more densely packed in the beginning, or maybe there was just less stuff.

Steady State Universe

Theories were proposed that suggested the universe was non changing in fundamental appearance.


Maybe the amount of matter was changing over time, so that the average density of the universe remained constant.

Big Bang

Another option was that the density was greater earlier in the history of the universe.

Also, the universe would have to be hotter too.

4.6 Big Bang Implications




The amount of He-3 present in the universe can be used to figure out the temperatures necessary in the very early universe. Then based on the expansions rates, we can figure out a temperature that the universe should be at now.

Rough estimates predicted 3-5 K.

If the universe were a blackbody, this should be convertible to a peak wavelength.

Blackbody radiation for a temperature of around 3 K.

5. Cosmic Microwave Background




Two scientists working at Bell Labs kept recording a signal that they thought was noise - and couldn't get rid of no matter what they tried. They called some colleagues who were involved in microwave astronomy.

The Holmdel Horn Antenna that Penzias and Wilson used to first measure the CMB.

By NASA - Great Images in NASA
Description, Public Domain,
<https://commons.wikimedia.org/w/index.php?curid=6463768>


5.7 Isotropy/Anisotropy

Is it the same everywhere?



Data from the first 9 Minutes of the COBE satellites mission.

[Astrophysical Journal, Part 2 - Letters](#)
 (ISSN 0004-637X), vol. 354, May 10,
 1990, p. L37-L40




The CMB has a dipole anisotropy. This is likely due to the motion of the local group of galaxies compared to the rest of the visible universe, or the Hubble Flow.

Dipole Anisotropy


COBE dipole data

How to make a CMB?



- Start with a really hot dense beginning: $T \gg 10^4 \text{ K}$. At these temperatures, baryonic matter will be completely ionized.
- Any photons will scatter off electrons and the universe will be opaque.
- A dense, hot, and opaque medium will produce blackbody radiation.

The ionized, opaque universe before recombination



- As the universe expands, the temperature cools. Around $T \sim 3000 \text{ K}$, the protons and free electrons combine to form neutral atoms.
- Now, the blackbody photons are free to stream throughout the universe.

Transparent universe after recombination

Cooling of the CMB

The energy density of a photon gas:

$$u = \left(\frac{4\sigma}{c} \right) T^4 \quad (8)$$

and the pressure:

$$P = \frac{u}{3} \quad (9)$$

From the first law of thermodynamics.

$$dQ = dE + PdV \quad (10)$$

Since the universe is homogeneous and isotropic, we expect there to be no heat flow: $dQ = 0$.

Thus, we get:

$$\frac{dE}{dt} = -P(t) \frac{dV}{dt} \quad (11)$$

Substituting u and P from above, and taking derivatives, and simplifying:

$$\frac{1}{T} \frac{dT}{dt} = -\frac{1}{3V} \frac{dV}{dt} \quad (12)$$

since the volume of the universe is proportional to the scale factor cubed:

$$V(t) \propto a(t)^3 \quad (13)$$

we can write:


$$\frac{1}{T} \frac{dT}{dt} = -\frac{1}{a} \frac{da}{dt} \quad (14)$$

which can be reexpressed as:

$$\frac{d}{dt}(\ln T) = -\frac{d}{dt}(\ln a) \quad (15)$$

which implies:

$$T(t) \propto a(t)^{-1} \quad (16)$$



[NASA / WMAP Science Team](#)

5.8 Olber's Paradox Resolution

The universe is not infinitely old.

Can't say how big it is, but only what we can see. (visible universe)