


Celestial Mechanics

1. [Basic Orbits](#)
 1. [Why circles?](#)
 2. [Tycho Brahe](#)
 3. [Johannus Kepler](#)
 4. [Kepler's 3 laws of orbiting bodies](#)
2. [Newtonian Mechanics](#)
3. [Newton's Laws](#)
 1. [Law of Gravitation](#)
 2. [The shell theorem.](#)
 3. [Work and Energy.](#)
4. [Potential Energy](#) →
 1. [Potential Plot](#)
 2. [Escape Speed](#)
5. [Kepler's Laws](#)
 1. [Cartesian & Polar Coordinates](#)
 2. [2nd Law](#)
 3. [1st Law - Ellipses](#)
6. [Third Law](#)
7. [Special Orbits](#)
 1. [Low Earth](#)
 2. [Geostationary](#)
 3. [Earth and Moon](#)
 4. [Hohmann Transfer Orbit](#)
8. [Lagrange Points](#)
 1. [Find L1](#)
9. [The Virial Theorem](#)
 1. [The Vis Viva equation](#)

I. Basic Orbits

I.1 Why circles?



The Copernican System consisted of circular orbits centered around the sun.


Copernicus was right: the earth and the other planets do orbit around the sun. However, he was still operating under the assumption that all the celestial bodies had to move in perfect circles. This turned out to be a rather major flaw in the framework.

Other astronomers of the era were unable to match the predictions of the Copernican system with observations. More 'tweaks' were added to the Copernican systems of circular heliocentric orbits in order to 'save the appearances'. In order to make the model match with observations, they still had to employ epicycles and so forth. However, the seed was planted.

The sun should be at the center!

I.2 Tycho Brahe

Late 16th century. Hybrid system: geoheliocentric model had the planets going around the sun, but the sun still went around the Earth. Close...



Danish astronomer Tycho Brahe [1546-1601]. Made many very good measurements of the stars and planets.

1.3 Johannes Kepler




German mathematician, astronomer, and astrologer. [1571-1630] Kepler was Tycho's assistant and was believed in the Copernican system.

Kepler played a major role in the 17th-century scientific revolution. He is best known for his laws of planetary motion, based on his works *Astronomia nova*, *Harmonices Mundi*, and *Epitome of Copernican Astronomy*. These works also provided one of the foundations for Newton's theory of universal gravitation.

1.4 Kepler's 3 laws of orbiting bodies

1. A planet orbits the sun in an ellipse. The Sun is at one focus of that ellipse.
2. A line connecting a planet to the Sun sweeps out equal areas in equal times
3. The square of a planet's orbital period is proportional to the cube of the average distance between the planet and the sun: $P^2 \propto a^3$.

Ellipses



An ellipse satisfies this equation:


$$r + r' = 2a \quad (1)$$

- a is the **semi-major axis**
- r and r' are the distances to the ellipse from the two **focal points**, F and F' .
- e represents the eccentricity of the ellipse ($0 \leq e \leq 1$)

The **semi-major axis** is one-half the length of the long (i.e. major) axis of the ellipse.


Conic Sections

The circle is one of several conic sections. (There's really nothing all that special about circles.)



See Apollonius - On Conics.

Elliptical Orbits




The first law for Kepler holds that the orbiting body (i.e. planet) will trace out an elliptical path as it orbits the central body (i.e. the sun). The central body is located at the **principle focus**. When the distance between the planet and the sun is the shortest, the planet is said to be at **perihelion**, or the point of closes approach. When that distance is greatest, the planet is located the **aphelion**.

Polar Coordinates

$$r = \frac{a(1 - e^2)}{1 + e \cos \theta} \quad (2)$$

The polar coordinate equation can be obtained by starting with the basic ellipse equation Eq. (1). Take the point at the top of the semi-minor axis, where $r = r'$. Since $r + r' = 2a$ we know that $r = a$. Using the Pythagorean theorem, we can say that:

$$r^2 = b^2 + a^2 e^2$$




Ellipse - the point where $r = r'$

Putting a in for r yields:

$$b^2 = a^2(1 - e^2) \quad (3)$$

Or, to have it more explicitly as the ratio of the two axes:

$$e = \sqrt{1 - \frac{b^2}{a^2}} \quad (4)$$



Ellipse - and polar coordinates r and θ .

For polar coordinates, we'll use r and θ , r being the distance from the principle focus and θ the angle measured counterclockwise from major axis of the ellipse. Again, from Pythagorus:


$$r'^2 = r^2 \sin^2 \theta + (2ae + r \cos \theta)^2$$

Expanding the 2nd RHS term:

$$\begin{aligned} r'^2 &= r^2 \sin^2 \theta + 4a^2e^2 + 4aer \cos \theta + r^2 \cos^2 \theta \\ &= r^2 + 4ae(ae + r \cos \theta) \end{aligned} \quad (5)$$


and using the fact that $r'^2 = (2a - r)^2$, we can obtain the polar equation for the ellipse shown above in eq (2).

Deviation from a Circles



Mars' orbit [$e = 0.0934$] looks a lot like a circle.

Equal Areas in Equal Times



The blue areas in these figures will be the same if $t_2 - t_1$ is the same.

Kepler's second law describes the areas swept out by an orbiting planet in a given time and says that for any given interval of time, the areas swept out will always be equal.

The other conics

Parabolas: $e = 1$

$$r = \frac{2p}{1 + \cos \theta} \quad (6)$$

Hyperbolas: $e > 1$

$$r = \frac{a(e^2 - 1)}{1 + e \cos \theta} \quad (7)$$

C/2023 E3 ZTF



Astronomy Picture of the Day

APOD of the comet

C/2023 E3 ZTF

Home / Tools / Small-Body Database Lookup

Small-Body Database Lookup

Search...

C/2022 E3 (ZTF)
 Classification: Hyperbolic Comet SPKID: 1003845 Related Links: [Ephemeris](#)

[Orbit Viewer](#) [show]

[Orbit Parameters](#) [hide]

Osculating Orbital Elements

Epoch 2459810.5 (2022-Aug-19.0) TDB
 Reference: [JPL 41](#) (heliocentric IAU76/J2000 ecliptic)

Element	Value	Uncertainty (1-sigma)	Units
e	1.000272119577545	5.2478E-7	
a	-4087.344214344506	7.8816	au
q	1.112246380888537	8.49E-7	au
i	109.1690286757335	1.4057E-5	deg
node	302.5543157083279	1.2995E-5	deg
peri	145.8152818804569	3.1658E-5	deg
M	-0.0005536428368530141	1.6014E-6	deg
tp	2459957.286944795938 2023-Jan-12.78694480	3.3302E-5	TDB
period	<i>not defined</i>	<i>not defined</i>	d
n	3.771744398813422E-6	1.091E-8	deg/d
Q	<i>not defined</i>	<i>not defined</i>	au

Lookup the comet in the small body database

https://ssd.jpl.nasa.gov/tools/sbdb_lookup.html/?des=2022%20E3

Small Body Database

2. Newtonian Mechanics

3. Newton's Laws

In words:

1. No change in velocity if there is no net force acting.
2. The net force acting on a body will cause a proportional acceleration inversely proportional to the mass of that body.
3. Forces come in equal and opposite pairs

In symbols:

1. $\Delta \mathbf{v} = \mathbf{0}$ if $\mathbf{F}_{\text{net}} = \mathbf{0}$
2. $\mathbf{F}_{\text{net}} = \sum_{i=1}^n \mathbf{F}_i = m\mathbf{a}$
3. $\mathbf{F}_{AB} = -\mathbf{F}_{BA}$

3.5 Law of Gravitation



Two masses and a gravitational interaction

The scalar form:

$$F = G \frac{Mm}{r^2} \quad (8)$$

And in vector form:

$$\mathbf{F}_{21} = -G \frac{Mm}{r^2} \hat{\mathbf{r}} \quad (9)$$

This is Newton's **Law of Universal Gravitation**. It considers two masses: M and m and the distance between them r . Also included is G , the **Universal Gravitational Constant**. ($G = 6.67430 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ From nist <http://physics.nist.gov/cgi-bin/cuu/Value?bg>). The direction of the force is always attractive and is along a line connecting the centers of the two masses.

\mathbf{F}_{21} is the force applied on object 2 by object 1.

r is the distance between the two objects ($|\mathbf{r}_{12}|^2$)

And $\hat{\mathbf{r}}_{12}$ is the unit vector pointing from object 1 to object to 2.

$$\hat{\mathbf{r}}_{12} = \frac{\mathbf{r}_2 - \mathbf{r}_1}{|\mathbf{r}_2 - \mathbf{r}_1|}$$




Cavendish

Example Problem
 #1:

3.6 The shell theorem.

Show that a spherical mass (of density $\rho(\mathbf{R})$) acts as a point mass for objects outside.



Consider a mass m a distance r away from the center of a spherical mass M . The radius of the spherical mass is R_0 . First, the force from the ring which a mass dM is:

$$dF_{\text{ring}} = G \frac{m dM_{\text{ring}}}{s^2} \cos \phi$$

The Shell Theorem

We can re-express the mass of the ring in terms of its density $\rho(R)$. (The density is at most only a function of the radius, i.e. spherically symmetric)

$$dM_{\text{ring}} = \rho(R) dV_{\text{ring}}$$

The volume of a ring is just its thickness: dR , times its width: $Rd\theta$, times its circumference: $2\pi R \sin \theta$ which equals:

$$dM_{\text{ring}} = \rho(R) 2\pi R^2 \sin \theta dR d\theta$$

The $\cos \phi$ can be expressed as:

$$\cos \phi = \frac{r - R \cos \theta}{s}$$

and s can be written as the following (from Pythagorus)

$$s = \sqrt{(r - R \cos \theta)^2 + R^2 \sin^2 \theta} = \sqrt{r^2 - 2rR \cos \theta + R^2}$$

Let's put all these back into the force from the ring:

$$\begin{aligned} dF_{\text{ring}} &= G \frac{m dM_{\text{ring}}}{s^2} \cos \phi \\ &= Gm [\rho(R) 2\pi R^2 \sin \theta dR d\theta] \times \left[\frac{1}{s^2} \right] \times \left[\frac{r - R \cos \theta}{s} \right] \end{aligned}$$

Then, we'll have to integrate over both variables. dR will range from 0 to R_0 and $d\theta$ from 0 to π to get all the rings that make up the entire sphere.

$$F = 2\pi Gm \left[\int_0^{R_0} \int_0^\pi \frac{\rho(R) (rR^2 \sin \theta - R^3 \sin \theta \cos \theta)}{(r^2 + R^2 - 2rR \cos \theta)^{3/2}} d\theta dR \right] \quad (10)$$

Integrating over θ yields:

$$F = \frac{Gm}{r^2} \int_0^{R_0} 4\pi R^2 \rho(R) dR \quad (11)$$

However, the mass of a shell of thickness dR is just:

$$dM_{\text{shell}} = 4\pi R^2 \rho(R) dR$$

which is just quantity inside the integrand. Thus the force from on shell will be,

$$dF_{\text{shell}} = \frac{Gm dM_{\text{shell}}}{r^2}$$

This force acts as if it is located at the center of the shell.

We can then integrate over all the mass shells and the above becomes:

$$\mathbf{F} = G \frac{mM}{r^2} \quad (12)$$

which is just the equation for the force between two point masses.


little g

$$\mathbf{F} = G \frac{M_{\oplus} m}{(R_{\oplus} + h)^2} \quad (13)$$

Let's consider the gravitational force a distance h above the surface of the earth. Since $\mathbf{F} = m\mathbf{a}$, we can see that the acceleration \mathbf{a} will be equal to:

$$\mathbf{a} = G \frac{M_{\oplus}}{R_{\oplus}^2} = g \quad (14)$$

We generally call this value 'little g '.




little g is not the same everywhere.

Example Calc [NB](#) →

gravity anomalies measured by GRACE

[NASA/JPL/University of Texas Center for Space Research](#)

3.7 Work and Energy



$$U_f - U_i = \Delta U = - \int_{\mathbf{r}_i}^{\mathbf{r}_f} \mathbf{F} \cdot d\mathbf{r} \quad (15)$$


The change in potential energy is given by the negative of the work done by gravity. Expressing it in the integral above allows for changing force as a function of position (in contrast to the elementary definition: $\mathbf{W} = \mathbf{F} \cdot \mathbf{d}$)

A point mass m moves from \mathbf{r}_i to \mathbf{r}_f in a gravitational field.

4.

Find the work done on an object by gravity as it moves from point \mathbf{P} to infinity, where the potential energy is defined as 0.

$$W = \int_R^{\infty} \mathbf{F}(\mathbf{r}) \cdot d\mathbf{r}$$



What is the potential energy at point P ?

The force is always against the displacement, so

$$\mathbf{F}(\mathbf{r}) \cdot d\mathbf{x} = -\frac{GM_E m}{r^2} dr$$

We can pull the constants out of the integral, and we are left with something very simple to integrate:

$$W = -GM_E m \int_R^\infty \frac{1}{r^2} dr = \left[\frac{GM_E m}{r} \right]_R^\infty$$

Evaluating at the limits of ∞ and R , we see that the work done is:

$$W = \left[\frac{GM_E m}{r} \right]_R^\infty = 0 - \frac{GM_E m}{R} = -\frac{GM_E m}{R}$$


From the original definition of work and potential energy:

$$\Delta U = U_f - U_i = -W$$

thus, since $U_\infty = 0$ we can say: $U = W$ And now we have a potential energy function for masses at a distance r from the center of the Earth:

$$U = -\frac{GM_e m}{r}$$

4.8 Potential Plot



The potential energy as a function of position for a body in a gravitational interaction.

4.9 Escape Speed

How fast to get something to reach infinity, (and stop there)?

$$E = \frac{1}{2}mv^2 - G\frac{Mm}{r} \tag{16}$$

This is the total energy of a particle: kinetic plus potential. Since the particle will be at $r = \infty$ and will have stopped then the total energy must be zero. However, due to conservation of energy, the total mechanical energy will be the same. Thus, $\frac{1}{2}mv^2 - G\frac{Mm}{r} = 0$ at all times. Or:

$$\frac{1}{2}mv^2 = G\frac{Mm}{r}$$



which leads to

$$v_{\text{escape}} = \sqrt{\frac{2GM}{r}} \quad (17)$$

For Earth, $v_{\text{escape}} = 11182.4$ m/s. Apollo 11 traveled at 10423 m/s [nasa ref]. It got a little help from the moons gravity too.

5. Kepler's Laws

5.10 Cartesian & Polar Coordinates



Polar Coordinates r and θ .

In polar coordinates, we have different unit vectors: $\hat{\theta}$ and \hat{r} .

$$\hat{r} = \hat{i} \cos \theta + \hat{j} \sin \theta \quad (18)$$

and

$$\hat{\theta} = -\hat{i} \sin \theta + \hat{j} \cos \theta \quad (19)$$

We can also express some derivatives:

$$\frac{d\hat{r}}{d\theta} = -\hat{i} \sin \theta + \hat{j} \cos \theta = \hat{\theta} \quad (20)$$

$$\frac{d\hat{\theta}}{d\theta} = -\hat{i} \cos \theta - \hat{j} \sin \theta = -\hat{r} \quad (21)$$

And using the chain rule, we can also express time derivatives:

$$\frac{d\hat{r}}{dt} = \frac{d\hat{r}}{d\theta} \frac{d\theta}{dt} = \hat{\theta} \frac{d\theta}{dt} \quad (22)$$

$$\frac{d\hat{\theta}}{dt} = \frac{d\hat{\theta}}{d\theta} \frac{d\theta}{dt} = -\hat{r} \frac{d\theta}{dt} \quad (23)$$

These will be useful later.

5.11 2nd Law

To prove the second law, we can show how Conservation of Angular Momentum results in the **equal areas in equal times** statement.

Angular Momentum

$$\vec{L} = \vec{r} \times \vec{p} \quad (24)$$

where $\vec{p} = m\vec{v}$

Take the time derivative of \vec{L}


$$\frac{d\vec{L}}{dt} = \frac{d\vec{r}}{dt} \times \vec{p} + \vec{r} \times \frac{d\vec{p}}{dt} = \vec{v} \times m\vec{v} + \vec{r} \times m\frac{d\vec{v}}{dt} \quad (25)$$

But, this is quickly seen to be zero for a central force:

$$\frac{d\vec{L}}{dt} = 0 \quad (26)$$

$$\vec{L} = \vec{r} \times m\vec{v} = mrv_t\hat{k} = L\hat{k} \quad (27)$$

The total area is the sum of the two triangles:



$$\Delta A \approx \frac{1}{2}r(v_t\Delta t) + \frac{1}{2}(v_r\Delta t)(v_t\Delta t) \quad (28)$$

Thus, in the limit of $r \gg v_r\Delta t$:

Motion's of an orbiter during a short time interval Δt

$$\Delta A \approx \frac{1}{2}r(v_t\Delta t) \quad (29)$$

Next, take the time derivative of the changing area:

$$\frac{dA}{dt} = \frac{1}{2}rv_t = \frac{1}{2} \frac{L}{m} \quad (30)$$

which shows the area in a given time is constant.

5.12 1st Law - Ellipses

The angular momentum per unit mass will be given by:

$$\frac{L}{m} = r^2 \frac{d\theta}{dt} \quad (31)$$

and the centrally directed gravitational force is given by:

$$\vec{F} = -\frac{GMm}{r^2}\hat{r} = m\frac{d\vec{v}}{dt} \quad (32)$$

The acceleration of the orbiting body is therefore:

$$\mathbf{a} = \frac{d\vec{v}}{dt} = -\frac{GM}{r^2}\hat{r} \quad (33)$$

but

$$\hat{r} = -\left(\frac{d\theta}{dt}\right)^{-1} \frac{d\hat{\theta}}{dt} \quad (34)$$

We can put eq (34) into the acceleration equation (33)

$$\frac{d\vec{v}}{dt} = \frac{GM}{r^2} \left(\frac{d\theta}{dt} \right)^{-1} \frac{d\hat{\theta}}{dt} \quad (35)$$


Combining this with (31)

$$\frac{L}{GMm} \frac{d\vec{v}}{dt} = \frac{d\hat{\theta}}{dt} \quad (36)$$

Integrating both sides of this equation yields:

$$\frac{L}{GMm} \vec{v} = \hat{\theta} + \vec{e} \quad (37)$$

The \vec{e} is a constant of integration that depends on the initial conditions, which we may choose.



We'll say that at $t = 0$, the orbiter is at perihelion and we can orient the axes so that its motion is entirely in the $+y$ direction, in other words: $\vec{e} = e\hat{j}$ where e is a constant. Now:

$$\frac{L}{GMm} \vec{v} = \hat{\theta} + e\hat{j} \quad (38)$$

We can take the dot product with $\hat{\theta}$ of both sides:

$$\frac{L}{GMm} \vec{v} \cdot \hat{\theta} = \hat{\theta} \cdot \hat{\theta} + e\hat{j} \cdot \hat{\theta} \quad (39)$$

The initial conditions of an orbiting body.

Simplifying, using the definitions above (i.e. (19)):

$$\frac{L}{GMm} v_t = 1 + e \cos \theta \quad (40)$$

(v_t) is the **tangential velocity** of the orbiter. From the definition of angular momentum: $\mathbf{L} = \mathbf{r} \times \mathbf{p}$

$$mrv_t = L \quad (41)$$

which when put back into (40), yields:

$$\frac{L^2}{GMm^2 r} = 1 + e \cos \theta \quad (42)$$

or solving for r :

$$r = \frac{L^2}{GMm^2 (1 + e \cos \theta)} \quad (43)$$

This is just the polar coordinates expression for a conic section.

6. Third Law

Special Case: Circular orbits

For example, for an object orbiting the earth, we can write:

$$G \frac{M_E m}{r^2} = m \frac{v^2}{r}$$

We can solve this for v , the speed of the orbiting object:

$$v = \sqrt{\frac{GM_E}{r}}$$

Now we can use the gravitational force to calculate the expected period:

$$v = \sqrt{\frac{GM_e}{r}} = \frac{2\pi r}{T}$$

which can be solved for T :

$$T = \frac{2\pi r^{3/2}}{\sqrt{GM_E}}$$

Kepler figured out a version of this relationship before Newton:

$$T \propto r^{3/2}$$


The period is proportional to the three-halves power of the orbital radius.

General Case (i.e. ellipses)

$$T^2 = \frac{4\pi^2}{G(M + m)} a^3 \quad (44)$$

7. Special Orbits

7.13 Low Earth



Low Earth Orbit


Most of our satellites are located in Low Earth Orbit. Roughly defined as less than 2000 km above the surface of the Earth.

Orbital Periods there range between 93 minutes and 127 minutes (for circular orbits)

[Calc Demo: →](#)

[Wayfinder: Link →](#)

7.14 Geostationary




GeoStationary

7.15 Earth and Moon



Earth and Moon to scale

7.16 Hohmann Transfer Orbit




The semi-major axis of the transfer orbit:

$$a_{\text{transfer}} = \frac{r_e + r_m}{2} \quad (45)$$


A typical Hohmann Transfer Orbit

8. Lagrange Points



The basic free body diagram showing the forces on a small body near two larger ones.

The basic FBD and L1 point



There are five Lagrange points. Some are interesting for technology applications.

All 5 Lagrange points

8.17 Find L1

The sum of forces acting on the earth is equal to its mass times the acceleration, which for an object moving in a circular orbit is centripetal and equal to v^2/R :

$$\Sigma F_{\text{on Earth}} = \frac{GMm_E}{R^2} = \frac{m_E v^2}{R}$$

Thus we can say:

$$\frac{GM}{R} = v^2$$

but, based on the relation between speed and period, T :

$$v = \frac{2\pi R}{T}$$

or

$$v^2 = \frac{4\pi^2 R^2}{T^2}$$

So we can rewrite as:

$$\frac{GM}{R} = \frac{4\pi^2 R^2}{T^2}$$

Rearranging yields Kepler's Third Law

$$\frac{GM}{R^3} = \frac{4\pi^2}{T^2} \quad (46)$$

Now consider the sum of forces on the satellite:

$$\Sigma F_{\text{on Sat}} = \frac{GMm_{\text{sat}}}{(R-r)^2} - \frac{Gm_E m_{\text{sat}}}{r^2} = \frac{m_{\text{sat}} v_{\text{sat}}^2}{(R-r)}$$

This we can simplify to, using Kepler's Third law for the satellite:

$$\frac{GM}{R-r} - \frac{Gm_E(r-R)}{r^2} = v_{\text{sat}}^2 = \frac{4\pi^2(R-r)^2}{T_{\text{sat}}^2}$$

or

$$\frac{GM}{(R-r)^3} - \frac{Gm_E}{r^2(R-r)} = \frac{4\pi^2}{T_{\text{sat}}^2} \quad (47)$$

Lastly, since we want their periods to be the same: $T_{\text{sat}} = T$

$$\frac{GM}{(R-r)^3} - \frac{Gm_E}{r^2(R-r)} = \frac{GM}{R^3} \quad (48)$$

If we put in the masses of the two major bodies, the Earth and Sun for example, we can calculate the value of r in terms of R . For our earth-sun system, the L1 point would be located at

$$r = 0.009969 \text{ AU}$$

or about 1/100 of the earth-sun distance. Changing signs in the sums of forces could allow to calculate the L2 and L3 positions as well.

9. The Virial Theorem

The **Virial theorem** shows that for a gravitationally bound system in equilibrium, the total energy is one-half the time averaged potential energy.

Consider the quantity Q .

$$Q \equiv \sum_i \mathbf{p}_i \cdot \mathbf{r}_i \quad (49)$$

where \mathbf{p}_i is the linear momentum and \mathbf{r}_i the position of some particle i . The product rule gives the following if we take the time derivative of the right hand side.

$$\frac{dQ}{dt} = \sum_i \left(\frac{d\mathbf{p}_i}{dt} \cdot \mathbf{r}_i + \mathbf{p}_i \cdot \frac{d\mathbf{r}_i}{dt} \right) \quad (50)$$

We could also say:

$$\frac{dQ}{dt} = \frac{d}{dt} \sum_i m_i \frac{d\mathbf{r}_i}{dt} \cdot \mathbf{r}_i = \frac{d}{dt} \sum_i \frac{1}{2} \frac{d}{dt} (m_i r_i^2) = \frac{1}{2} \frac{d^2 I}{dt^2} \quad (51)$$

where I is the moment of inertia of the total number of particles:

$$I = \sum_i m_i r_i^2 \quad (52)$$

$$\frac{1}{2} \frac{d^2 I}{dt^2} - \sum_i \mathbf{p}_i \cdot \frac{d\mathbf{r}_i}{dt} = \sum_i \frac{d\mathbf{p}_i}{dt} \cdot \mathbf{r}_i \quad (53)$$

Now,

$$-\sum_i \mathbf{p}_i \cdot \frac{d\mathbf{r}_i}{dt} = -\sum_i m_i \mathbf{v}_i \cdot \mathbf{v}_i = -2 \sum_i \frac{1}{2} m_i v_i^2 = -2K \quad (54)$$

Thus:

$$\frac{1}{2} \frac{d^2 I}{dt^2} - 2K = \sum_i \mathbf{F}_i \cdot \mathbf{r}_i \quad (55)$$

where we have used the 2nd law of Newton ($\mathbf{F} = \frac{d\mathbf{p}}{dt}$)

Let \mathbf{F}_{ij} represent the force of interaction between two particles in the system. The force on i due to j . If you have a particle i , then we need to add up the force contributions from every other particle j where $j \neq i$.

$$\sum_i \mathbf{F}_i \cdot \mathbf{r}_i = \sum_i \left(\sum_{\substack{j \\ j \neq i}} \mathbf{F}_{ij} \right) \cdot \mathbf{r}_i \quad (56)$$

We can re-write the position of the i th particle as:

$$\mathbf{r}_i = \frac{1}{2}(\mathbf{r}_i + \mathbf{r}_j) + \frac{1}{2}(\mathbf{r}_i - \mathbf{r}_j) \quad (57)$$

which leads to

$$\sum_i \mathbf{F}_i \cdot \mathbf{r}_i = \frac{1}{2} \sum_i \left(\sum_{\substack{j \\ j \neq i}} \mathbf{F}_{ij} \right) \cdot (\mathbf{r}_i + \mathbf{r}_j) + \frac{1}{2} \sum_i \left(\sum_{\substack{j \\ j \neq i}} \mathbf{F}_{ij} \right) \cdot (\mathbf{r}_i - \mathbf{r}_j) \quad (58)$$

Since the 3rd law of Newton says: $\mathbf{F}_{ij} = -\mathbf{F}_{ji}$, the first term on the R.H.S. in the above equation will be zero. Thus:

$$\sum_i \mathbf{F}_i \cdot \mathbf{r}_i = \frac{1}{2} \sum_i \sum_{\substack{j \\ j \neq i}} \mathbf{F}_{ij} \cdot (\mathbf{r}_i - \mathbf{r}_j) \quad (59)$$

Assuming only gravitational interactions:

$$\mathbf{F}_{ij} = G \frac{m_i m_j}{r_{ij}^2} \hat{\mathbf{r}}_{ij} \quad (60)$$

we can write

$$\sum_i \mathbf{F}_i \cdot \mathbf{r}_i = -\frac{1}{2} \sum_i \sum_{\substack{j \\ j \neq i}} G \frac{m_i m_j}{r_{ij}^3} (\mathbf{r}_j - \mathbf{r}_i)^2 \quad (61)$$

$$- \frac{1}{2} \sum_i \sum_{\substack{j \\ j \neq i}} G \frac{m_i m_j}{r_{ij}} \quad (62)$$

The potential energy U_{ij} between the i and j particles is just the term in the sums:

$$U_{ij} = -G \frac{m_i m_j}{r_{ij}} \quad (63)$$

Thus,

$$\sum_i \mathbf{F}_i \cdot \mathbf{r}_i = \frac{1}{2} \sum_i \sum_{\substack{j \\ j \neq i}} U_{ij} = U \quad (64)$$

Now we can fill out eq: (55)

$$\frac{1}{2} \left\langle \frac{d^2 I}{dt^2} \right\rangle - 2 \langle K \rangle = \langle U \rangle \quad (65)$$

where we have replaced all the quantities with the time averaged values. The average of $\frac{d^2 I}{dt^2}$ is going to be:

$$\left\langle \frac{d^2 I}{dt^2} \right\rangle = \frac{1}{\tau} \int_0^\tau \frac{d^2 I}{dt^2} dt \quad (66)$$

$$= \frac{1}{\tau} \left(\frac{dI}{dt} \Big|_\tau - \frac{dI}{dt} \Big|_0 \right) \quad (67)$$

but for periodic motions like orbits:

$$\frac{dI}{dt} \Big|_\tau = \frac{dI}{dt} \Big|_0 \quad (68)$$

So

$$\left\langle \frac{d^2 I}{dt^2} \right\rangle = 0 \quad (69)$$

which leads finally to

$$-2 \langle K \rangle = \langle U \rangle$$

or in terms of the total energy E

$$\langle E \rangle = \frac{1}{2} \langle U \rangle$$

The Virial Theorem

$$-2 \langle K \rangle = \langle U \rangle$$

or in terms of the total energy E

$$\langle E \rangle = \frac{1}{2} \langle U \rangle$$

The Virial Theorem: Uses

It allows for statistical results for many body systems.

For example, with galaxies: Since $K = \frac{1}{2} M v^2$ and $U = \frac{GM^2}{R}$, we can obtain:

$$M = \frac{v^2 R}{G}$$

which relates two observable values, v and R , with a non observable, but very interesting value, the mass of the galaxy.

9.18 The Vis Viva equation

$$v^2 = GM \left(\frac{2}{r} - \frac{1}{a} \right)$$